

Incident-energy and system-size dependence of directed flow

Gang Wang (for the STAR[‡] Collaboration)

University of California, Los Angeles, California 90095, USA

E-mail: gwang@physics.ucla.edu

Abstract. We present STAR's measurements of directed flow for charged hadrons in Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV and 62.4 GeV, as a function of pseudorapidity, transverse momentum and centrality. We find that directed flow depends on the incident energy, but not on the system size. We extend the validity of limiting fragmentation hypothesis to different collision systems.

1. Motivation

Directed flow is quantified by the first harmonic (v_1) in the Fourier expansion of the azimuthal distribution of produced particles with respect to the reaction plane [1]. It describes the collective sideward motion of produced particles and nuclear fragments, and carries information from the very early stage of the collision [2]. Charged-hadron v_1 results at RHIC have been previously reported over a wide range of incident energies in Au+Au collisions [3, 4, 5, 6], and they support the limiting fragmentation hypothesis [4, 5, 7]. In RHIC run V (2005), lighter nuclei were collided at $\sqrt{s_{NN}} = 200$ GeV and 62.4 GeV, which enables us to study the system-size dependence of directed flow, as well as the incident-energy dependence.

2. Data sets and the study

This study is based on eight million 200 GeV Au+Au events, five million 62.4 GeV Au+Au events, twelve million 200 GeV Cu+Cu events, and eight million 62.4 GeV Cu+Cu events. All of them were obtained from a minimum-bias trigger. All errors are statistical. Tracks of charged particles are reconstructed by STAR's main TPC [8] and forward TPCs [9], with the pseudorapidity (η) coverage of $|\eta| < 1.3$ and $2.5 < |\eta| < 4.0$, respectively. The centrality definition and cuts used are the same as in Ref. [10].

At RHIC energies, it is a challenge to measure v_1 accurately due to the small signal of v_1 itself and the large systematic error arising from non-flow correlations. In this talk, STAR's preliminary $v_1\{\text{ZDC-SMD}\}$ results are obtained with non-flow effects

[‡] For the full list of STAR authors and acknowledgements, see appendix 'Collaborations' in this volume.

Table 1. The resolution of the 1st-order full event plane provided by STAR ZDC-SMDs, as determined from the sub-event correlation between east and west SMDs. The errors in the table are statistical.

Centrality	200 GeV Au+Au	62.4 GeV Au+Au	200 GeV Cu+Cu	62.4 GeV Cu+Cu
70% – 80%	0.296 ± 0.003	0.179 ± 0.005		
60% – 70%	0.348 ± 0.003	0.185 ± 0.004		
50% – 60%	0.382 ± 0.002	0.176 ± 0.005	0.140 ± 0.003	0.039 ± 0.011
40% – 50%	0.397 ± 0.002	0.167 ± 0.005	0.144 ± 0.003	0.043 ± 0.011
30% – 40%	0.390 ± 0.002	0.138 ± 0.006	0.147 ± 0.003	0.032 ± 0.015
20% – 30%	0.365 ± 0.002	0.110 ± 0.008	0.121 ± 0.004	
10% – 20%	0.309 ± 0.003	0.081 ± 0.010	0.095 ± 0.005	
5% – 10%	0.220 ± 0.006		0.059 ± 0.008	
0 – 5%	0.120 ± 0.002		0.059 ± 0.008	

minimized by determining the event plane from spectator neutrons, the same approach as used in Ref. [5, 6]. The resolution, as defined in Ref. [1], of the first-order event plane reconstructed with STAR ZDC-SMDs [11] is listed in Table 1. The magnitude of the event plane resolution increases with the spectator v_1 and the detector efficiency of the ZDC-SMDs, and the latter is smaller for a lower incident energy and/or a smaller collision system. The method $v_1\{\text{ZDC-SMD}\}$ fails in some centralities, where the event plane resolution is consistent with zero.

3. Results

Figures in this talk follow the same convention as in Ref. [5].

Fig. 1 shows v_1 as a function of pseudorapidity, η , in 30% – 60% most central Au+Au and Cu+Cu collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV and 64.2 GeV. Data points in Fig. 1 fall into two bands, and the higher band corresponds to the lower energy, which reveals the incident-energy dependence of $v_1(\eta)$. For the pseudorapidity and centrality range we studied, $v_1(\eta)$ in Au+Au collisions and Cu+Cu collisions are consistent within errors if both of them were obtained at the same incident energy. It is interesting that v_1 does not change from Au+Au collisions to Cu+Cu collisions, while the system size is reduced by 1/3. A good consistency of v_1 between Au+Au and Cu+Cu collisions is observed even for the region near midrapidity, where v_2 in Cu+Cu collisions is considerably lower than that in Au+Au collisions [12]. Unlike v_2/ϵ which scales with $\frac{1}{S} \frac{dN_{ch}}{dy}$ (interpreted to be the participant density [12] or the system length [13]), v_1 is found to be independent of the system size. Instead, it scales with the incident energy. A possible explanation to the different scalings for v_2/ϵ and v_1 might come from the way in which they are developed : to produce v_2 , intensive momentum exchanges among particles are needed (and the number of momentum exchanges is related to the participant density or the system length), while to produce v_1 , one in principle needs only different rapidity losses (related to the incident energy) for particles with different distances to the center of the




Figure 1. Charged-hadron v_1 vs. η , in Au+Au and Cu+Cu collisions at 200 GeV and 62.4 GeV. The plotted errors are statistical.




Figure 2. Charged-hadron v_1 vs. $\eta - y_{\text{beam}}$, in Au+Au and Cu+Cu collisions at 200 GeV and 62.4 GeV. The plotted errors are statistical.

participant zone.

In Fig. 2, we re-arrange the data points of Fig. 1 in the projectile frame, where zero on the horizontal axis corresponds to the beam rapidity, y_{beam} , for each of the incident energies. Within three units from y_{beam} , most data points fall into a universal curve of v_1 as a function of $\eta - y_{\text{beam}}$. This incident-energy scaling behavior of directed flow suggests that the limiting fragmentation hypothesis [7] holds even for different collision systems.

The transverse-momentum (p_t) dependence of v_1 for charged hadrons is shown in Fig. 3. The upper panel shows the results measured in the main TPC, and the lower panel, in the forward TPCs. Since $v_1(\eta, p_t)$ is asymmetric about $\eta = 0$, the integral of $v_1(\eta, p_t)$ over a symmetric η range goes to zero. We change $v_1(\eta, p_t)$ of particles with negative η into $-v_1(-\eta, p_t)$, and integrate over all η . At 200 GeV, $v_1(p_t)$ in Au+Au and Cu+Cu collisions are very close to each other (better visible in the forward pseudorapidity region), and $v_1(p_t)$ saturates above $p_t \approx 0.8$ GeV/ c . At 62.4 GeV, $v_1(p_t)$ reaches its maximum in magnitude at $p_t \approx 0.6$ GeV/ c , then saturates ($|\eta| < 1.3$) or slightly decreases ($2.5 < |\eta| < 4.0$). Within statistical errors, the incident-energy scaling holds in $v_1(p_t)$ as well.

Fig. 4 shows charged-hadron p_t -integrated v_1 as a function of centrality for the two collision systems at 200 GeV and 62.4 GeV, and the results are presented in two panels defined in the same way as Fig. 3. On the whole, the magnitude of integrated v_1 approaches zero in central collisions and increases as the collisions become more peripheral. At 200 GeV, the p_t -integrated v_1 in Au+Au collisions is consistent with that in Cu+Cu collisions in almost all centrality bins, in regions both near the midrapidity ($|\eta| < 1.3$) and away from it ($2.5 < |\eta| < 4.0$). This system-size independence is seemingly true at 62.4 GeV. We can see that the incident-energy scaling of v_1 works well as a function of centrality, instead of the participant density, with which the v_2/ϵ scales [12]. This indicates that the initial geometrical shape, but not the density of produced particles, plays an important role in the formation of directed flow.




Figure 3. Charged-hadron v_1 vs. p_t , in 30%–60% Au+Au and Cu+Cu collisions at 200 GeV and 62.4 GeV. The upper panel shows the results measured in the main TPC, and the lower panel, in the forward TPCs. The plotted errors are statistical.




Figure 4. Charged-hadron p_t -integrated v_1 vs. centrality, in Au+Au and Cu+Cu collisions at 200 GeV and 62.4 GeV. The upper panel shows the results measured in the main TPC, and the lower panel, in the forward TPCs. The plotted errors are statistical.

References

- [1] Poskanzer A M and Voloshin S A 1998 *Phys. Rev. C* **58** 1671
- [2] Adams J *et al* (STAR Collaboration) 2005 *Nucl. Phys. A* **757** 102, and refs. therein
- [3] Adams J *et al* (STAR Collaboration) 2004 *Phys. Rev. Lett.* **92** 062301
- [4] Back B B *et al* (PHOBOS Collaboration) 2004 *J. Phys. G* **30** S1243; Back B B *et al* (PHOBOS Collaboration) 2006 *Phys. Rev. Lett.* **97** 012301
- [5] Adams J *et al* (STAR Collaboration) 2006 *Phys. Rev. C* **73** 034903
- [6] Wang G *et al* (STAR Collaboration) 2006 *Nucl. Phys. A* **774** 515
- [7] Benecke J, Chou T T, Yang C -N and Yen E 1969 *Phys. Rev.* **188** 2159
- [8] Anderson M *et al* 2003 *Nucl. Instr. Meth. A* **499** 659
- [9] Ackermann K H *et al* 2003 *Nucl. Instr. Meth. A* **499** 713
- [10] Adams J *et al* (STAR Collaboration) 2005 *Phys. Rev. C* **72** 014904
- [11] Adler C *et al* 2001 *Nucl. Instr. Meth. A* **470** 488; The STAR ZDC-SMD has the same structure as the STAR EEMC SMD: Allgower C E *et al* 2003 *Nucl. Instr. Meth. A* **499** 740; Crawford H *et al* 2003 STAR ZDC-SMD proposal *STAR Note SN-0448*
- [12] Back B B *et al* (PHOBOS Collaboration) 2006 *Preprint* nucl-ex/0610037; Voloshin S A *et al* (STAR Collaboration) these proceedings
- [13] Tang A H these proceedings